



## TABLE VARIETY

- *Excellent productivity.*
- *Medium early variety.*
- *Big oval to long oval tubers.*
- *White flesh.*

## PLANT & TUBER

|                  |                   |
|------------------|-------------------|
| FOLIAGE MATURITY | MEDIUM EARLY      |
| EMERGENCE        | MODERATE          |
| PLANT VIGOUR     | HIGH              |
| SKIN COLOUR      | YELLOW            |
| FLESH COLOUR     | WHITE             |
| TUBER SHAPE      | OVAL TO LONG OVAL |
| SHAPE UNIFORMITY | GOOD              |
| EYE DEPTH        | SHALLOW           |
| TUBER SIZE       | LARGE             |

## PESTS & DISEASES

|   |            |
|---|------------|
| RESISTANCE TO FOLIAGE LATE BLIGHT         | FAIRLY LOW |
| RESISTANCE TO TUBER LATE BLIGHT           | FAIRLY LOW |
| RESISTANCE TO PITTED COMMON SCAB          | MODERATE   |
| RESISTANCE TO NETTED COMMON SCAB          | MODERATE   |
| RESISTANCE TO PVY                         | FAIRLY LOW |
| RESISTANCE TO PCN G.rostochiensis (Ro1-4) | LOW        |
| RESISTANCE TO BLACK WART                  | HIGH       |

## QUALITY & USE

|                         |             |
|-------------------------|-------------|
| YIELD                   | VERY HIGH   |
| DRY MATTER CONTENT      | LOW         |
| COOKING TYPE            | A           |
| DISINTEGRATION          | VERY SLIGHT |
| DARKENING AFTER COOKING | NONE        |
| FRY COLOUR at harvest   | MEDIUM      |
| STORAGE ABILITY         | FAIRLY HIGH |

## PHYSIOLOGY

|                                  |             |
|----------------------------------|-------------|
| DORMANCY                         | MEDIUM      |
| RESISTANCE TO EXTERNAL DAMAGE    | FAIRLY HIGH |
| RESISTANCE TO INTERNAL DAMAGE    | FAIRLY HIGH |
| RESISTANCE TO INTERNAL RUST SPOT | FAIRLY HIGH |
| RESISTANCE TO DESPROUTING        | HIGH        |